



September, 2010

Volume 5, Issue 9

Thornton Natural Healthcare's Better Health News

www.thornton-health.com

Special Interest Articles:

- Bowel flora
- Vitamin A
- Supplementation and inflammatory bowel disease
- Diet and inflammatory bowel disease
- GAGs and inflammatory bowel disease
- Vitamin C
- Risks for inflammatory bowel disease

Inflammatory Bowel Disease

Inflammatory bowel disease exists when the intestines become chronically inflamed. Two manifestations of inflammatory bowel disease are Crohn's disease and ulcerative colitis. Symptoms of inflammatory bowel disease include abdominal cramps, pain diarrhea, weight loss and bleeding from the intestines. About 600,000 Americans have inflammatory bowel disease. Ulcerative colitis affects the interior lining of the colon and rectum. It can result in numerous health complications including rectal bleeding, diarrhea,

dehydration and increased risks of developing cancer. Crohn's disease usually causes ulcers along the length of the small and large intestines.. Complications from Crohn's disease can include obstruction of the intestine due to swelling and the formation of scar tissue, malnutrition, the development of fissures (small cuts or tears in the anal canal), abscesses (localized infection or collection of pus), and fistulas (an abnormal tunnel that forms between two structures of the body).

B₆ and Inflammatory Bowel Disease

Some of vitamin deficiencies may contribute to the severity of inflammatory bowel disease. The disease creates vitamin deficiency, which in turn makes the disease more severe. A study involving 30 male and 31 female patients with inflammatory bowel disease was published in the *American Journal of Gastroenterology* (2003;98 (1):112-117). It found that vitamin B₆ levels were significantly lower in patients with inflammatory bowel disease than they were in healthy controls. Furthermore,

vitamin B₆ levels were lower in patients who were experiencing a flare up in their symptoms than they were for patients in remission. Low vitamin B₆ levels were also associated with higher levels of what are known as inflammatory markers (these are chemicals that, when present, indicate inflammation) C-reactive protein is an example of an inflammatory marker; it was increased in patients with low B₆ levels.

Bowel Flora and Inflammatory Bowel Disease

Patients with inflammatory bowel disease had much higher concentrations of bacteria invading the intestinal lining than did the healthy subjects.

Normal bowel flora may play a role in inflammatory bowel disease (Crohn's disease and ulcerative colitis). An article appearing in the *Scandinavian Journal of Gastroenterology* (2001;36(Suppl 234):29-40) discussed the role of bowel flora and intestinal permeability in this disease. The intestinal tract is a long tube that goes from the mouth to the anus. The food you eat and everything inside the intestinal tract is still technically outside of the body. The cells lining the intestine form a barrier that selects what gets inside of the body and what stays outside of the body. If these cells fail to act as an effective barrier, too much intestinal permeability become a problem (sometimes called "leaky gut"). This article states that increased intestinal permeability leads to a lack of tolerance to the bacteria that normally exist in the bowel. There is evidence of the immune system working against the bacteria that exist in the bowel, perhaps even overreacting. Also, the makeup of the bacteria is different in these patients.

In patients with inflammatory bowel disease, the bacteria tend to invade the mucosa (lining of the

intestine). Research appearing in *Gastroenterology* (January 2002;122(1):44-54) compared colonoscopic biopsies in 305 patients with 40 normal controls. Patients with inflammatory bowel disease had much higher concentrations of bacteria invading the intestinal lining than did the healthy subjects. A normal, healthy intestinal lining has the ability to act as a barrier to bowel bacteria; this ability has been compromised in patients with inflammatory bowel disease.

The *Scandinavian Journal of Gastroenterology* article mentions that taking probiotics is sometimes helpful; patients with Crohn's disease tend to have less Bifidobacteria, and more Bacteroides, Eubacteria and Peptostreptococcus. Probiotics are supplements composed of living bacteria that, when taken, can be beneficial to the digestive tract. Generally these are lactic acid bacteria, like Lactobacilli, Bifidobacteria (the species found to be in short supply in patients with Crohn's disease), and Streptococci. These bacteria can release antibiotic-like substances, reduce the pH in the intestine, improve the absorption of nutrients and help stimulate the immune system.

Vitamin A and Inflammatory Bowel Disease

Vitamin A is among the nutrients that are deficient in patients with inflammatory bowel disease (ulcerative colitis or Crohn's disease). In a study appearing in *Hepato-Gastroenterology* (1991;38:391-395) compared 32 patients with inflammatory bowel disease (17 patients with Crohn's disease and 15 patients with ulcerative colitis) and compared them to healthy controls. The retinol (vitamin A) levels and

retinol-binding protein levels were lower in the patients with inflammatory bowel disease than in healthy controls. Ulcerative colitis patients who were successfully treated had their vitamin A level increase to normal, even without vitamin A supplementation. Similarly, vitamin A levels normalize in patients with Crohn's disease when the disease is inactive.

Inflammatory Bowel Disease & Supplementation

Inflammatory bowel disease is a general term to describe recurring inflammation of the small or large intestine. Inflammation often leads to ulceration and eventually scar tissue can form. This includes Crohn's disease and ulcerative colitis.

Patients with inflammatory bowel disease are often deficient in many nutrients. They commonly have problems absorbing nutrients, have poor appetites, are often on restrictive diets, or are nutrient deficient as a result of drug therapy. According to a review of research appearing in the *Annual Review of Nutrition* (1985;5:463-484), nutritional deficiencies are common in patients hospitalized with inflammatory bowel disease. Iron deficiency was found in 40% of the patients, 48% were deficient in vitamin B₁₂, between 54 and 64% were deficient in folate, between 14 and 33% were magnesium deficient, between 6% and 14% had a potassium deficiency, 21% were deficient in vitamin A, 12% were deficient in vitamin C, between 25% and 65% were deficient in 25-hydroxyvitamin D and between 40% and 50% had a zinc deficiency. Other vitamin deficiencies included vitamin K, copper and vitamin E.

According to a study of 114 patients with inflammatory bowel disease, appearing in the *Scandinavian Journal of Gastroenterology* (1979;14:1019-1024) found low serum folate in 59% of patients with chronic inflammatory bowel disease. There is an increased risk for colorectal cancer in patients with inflammatory bowel disease. Research appearing *Inflammatory Bowel Diseases* (2008 Feb;14(2):242-8), folic acid deficiency was associated with an increased risk

of colorectal cancer. Patients who had inflammatory bowel disease and both a folic acid deficiency and high homocysteine had 17 times as many cancerous lesions as patients who were not folic acid deficient.

Some of the vitamin deficiencies may contribute to the severity of the disease. The disease creates vitamin deficiency, which in turn makes the disease more severe. A study involving 30 male and 31 female patients with inflammatory bowel disease was published in the *American Journal of Gastroenterology* (2003;98(1):112-117). It found that vitamin B₆ levels were significantly lower in patients with inflammatory bowel disease than they were in healthy controls. Furthermore, vitamin B₆ levels were lower in patients who were experiencing a flare up in their symptoms than they were for patients in remission. Low vitamin B₆ levels were also associated with higher levels of what are known as inflammatory markers (these are chemicals that, when present, indicate inflammation) C-reactive protein is an example of an inflammatory marker; it was increased in patients with low B₆ levels.

Vitamin supplementation for patients with inflammatory bowel disease is sometimes a problem. For some patients, taking vitamin pills can exacerbate the condition. There are liquid vitamins that may be better tolerated (folic acid is not stable in a liquid multiple vitamin and should be given separately; several companies make a liquid folic acid supplement). Sometimes giving vitamins intravenously is a solution.

Some of the vitamin deficiencies may contribute to the severity of the disease. The disease creates vitamin deficiency, which in turn makes the disease more severe.

Diet and Inflammatory Bowel Disease

Diet can play a role in inflammatory bowel disease (ulcerative colitis and Crohn's disease). Food sensitivity may play a role in some cases. A small study published in *South African Family Practice* (June, 1995;16:393-399) resulted in six of seven patients with inflammatory bowel disease became symptom-free after following an elimination diet. Foods found to trigger reactions in various patients included pineapple, banana, cheese, tomato, coffee and milk.

In a study that appeared in *Hepato-Gastroenterology* (1990;37:72-80) looked at the effect an elemental diet had on patients with Crohn's disease. The elemental diet involves consuming all nutrients in a liquid form via a gastric feeding tube or IV, gives the inflamed gastrointestinal tract an opportunity to heal. With no food being consumed, digestion does not have to occur, giving the GI tract opportunity to reprieve from the very processes that inflame it. In this study, patients on an elemental diet actually improved more than patients who were given drug therapy.

Consumption of refined carbohydrates can aggravate symptoms. The *European Journal of Gastroenterology and Hepatology* (January, 1995;7(1):47-51) conducted a prospective, epidemiological study of 104 patients and found that patients with Crohn's disease and ulcerative colitis have a high intake of starch and sugar. Other research that appeared in *Epidemiology* (January 1992;3(1):47-52), followed the dietary habits of 142

patients with ulcerative colitis, 152 patients with Crohn's disease and 305 healthy controls over a period of three years. The relative risk of Crohn's disease increased with sugar intake and decreased with fiber intake. Eating fast food increased the risk for both Crohn's disease and for ulcerative colitis. These studies may support Elaine Gottschall's model of inflammatory bowel disease.

In her book, *Ending the Vicious Cycle*, Gottschall describes a possible mechanism for inflammatory bowel disease and a dietary solution. She reasons that we eat too many complex carbohydrates (disaccharides like table sugar and lactose in milk, and starches like bread, potatoes, rice and beans). Our own enzymes are not adequate enough to digest the large amounts of sugar and starch that we consume, so it passes into the intestine without being digested. The undigested food allows bacteria to grow, irritating the intestinal lining. The irritation of the intestine causes a further reduction in the amount of enzymes produced, resulting in more undigested food. Gottschall suggests that completely avoiding starches and sugar (the sugar that is in fresh fruit is permissible). Some patients, but not all, go into remission after following Gottschall's recommendations.

GAGs and Inflammatory Bowel Disease

The term GAGS is a shortened way of saying glycosaminoglycans, which are long molecules composed of sugar molecules (if a sugar molecule is a train car, GAGS are the train). Most of the GAGS in the body are used to form Proteoglycans, which are very important to structure. Proteoglycans are very important for the formation of cartilage. Perhaps the best-know proteoglycan is chondroitin sulfate. These molecules are not only important for joint cartilage, they help give form and structure to all areas of the body--including the intestines. Proteoglycans help form what is known as the basement membrane. The basement membrane acts as an

anchor to the epithelium of the intestine. The epithelium are the cells that provide the functionality of digestion; the basement membrane attaches to cartilage and to the epithelial cells and helps give the intestine form and structure. An article appearing in *The Lancet* (March 20, 1993;341:730- 731) discusses the fact that the glycosaminoglycans are altered in chronic inflammatory bowel disease. Bacteria, viruses, antigens and oxidative stress challenge the structural integrity of the intestine, glycosaminoglycans help protect against these challenges and help reduce intestinal permeability (leaky gut).

The Lancet
(March 20,
1993;341:730
- 731)
discusses the
fact that the
GAGs are
altered in
chronic
inflammatory
bowel
disease.

Vitamin C and Inflammatory Bowel Disease

You may have come across terms like “antioxidant”, “free radical” or “oxidative stress”. A free radical is a chemical that readily gives up an electron—it sort of fires the electron like a chemical bullet. Free radicals can damage tissue and produce inflammation; they create oxidation. Antioxidants are vitamins, nutrients and phytochemicals that act like little “bullet-proof vests” and help protect the cells.

One of the roles vitamin C has is as an antioxidant, protecting cells from oxidative stress. According to research

appearing in *Free Radical Research* (1995;22(2):131-143), patients with inflammatory bowel disease (Crohn's disease or ulcerative colitis) have reduced levels of vitamin C in the intestinal lining. Patients with Crohn's disease have levels that are 35% below normal and patients with ulcerative colitis have levels that are 73% below normal. The researchers state that most of the loss of vitamin C is due to oxidative stress from inflammatory cells. The loss of vitamin C makes the cells lining the intestines more vulnerable to oxidative stress.

Testimonial of the month

*If we knew what
we were doing,
we wouldn't call it
"research"--
Einstein*

I remember thinking that no one believed I was sick and accepting that I was going to feel bad forever. I have hypothyroidism and had gained weight, had zero energy, was freezing all the time, and felt like my mind was constantly in a fog. I had visited my family doctor who performed a blood test and told me my thyroid level was normal.

Then one day my throat began to feel like it was swelling shut. When I visited our school nurse she informed me my thyroid was swollen. After researching thyroid problems, I found I had many symptoms of hypothyroidism as I had suspected earlier. I found out I have an extensive family history of thyroid problems and requested to see a specialist. This endocrinologist listened to my symptoms and immediately told me I was depressed. I had to push him to take blood tests and ultrasound my thyroid. The ultrasound revealed an enlarged thyroid and many cysts, but he was not concerned and did not think I needed a follow-up. I received my blood test results in the mail and although they were in the normal range one test was high normal and the other low normal, another indication of hypothyroidism.

I requested to visit another specialist who treated me as though I were jumping from doctor to doctor in search of pills. This specialist stated he/she would give me medicine if my levels were 0.5 pts. higher and wanted me to come back in three months for another test to see if my levels had went up. By this time I had felt terrible for ten months and was giving up on finding anyone who would help me. Then a friend suggested I see Dr. Thornton. I was already a chiropractic patient of Dr. Thornton and knew him to be a compassionate person. I took the test results from the first specialist, told him my story, and walked out that day with a natural supplement for thyroid problems.

Within a month my levels were normal, I had lost twenty pounds and felt wonderful. I am able to teach and have conversations with parents and colleagues without struggling to keep my thoughts straight. Dr. Thornton is a blessing! Thank you!

—Z.S. from Aldrich, MO

Come visit our website!

www.thornton-health.com

Also visit us on [Facebook!](#)

205 South St. Stockton, MO
(417) 276-6306